arXiv:1209.3147v2 [gr-qc] 3 Oct 2012

Interacting generalized ghost dark energy in non-flat universe

Esmaeil Ebrahimi[1,3] * and Ahmad Sheykhi[2,3] †
[1] *Department of Physics, Shahid Bahonar University, PO Box 76175, Kerman, Iran*
[2] *Physics Department and Biruni Observatory, College of Sciences, Shiraz University, Shiraz 71454, Iran*
[3] *Research Institute for Astronomy and Astrophysics of Maragha (RIAAM), P. O. Box 55134-441, Maragha, Iran*

We investigate the generalized QCD ghost model of dark energy in the framework of Einstein gravity. First, we study the non-interacting generalized ghost dark energy in a flat FRW background. We obtain the equation of state parameter, $w_D = p/\rho$, the deceleration parameter, and the evolution equation of the generalized ghost dark energy. We find that, in this case, w_D cannot cross the phantom line ($w_D > -1$) and eventually the universe approaches a de-Sitter phase of expansion ($w_D \to -1$). Then, we extend the study to the interacting ghost dark energy in both flat and non-flat FRW universe. We find out that the equation of state parameter of the interacting generalized ghost dark energy can cross the phantom line ($w_D < -1$) provided the parameters of the model are chosen suitably. We also discuss the values of the deceleration parameter in all cases and present a numerical discussion of this parameter in different phases of the universe expansion.

I. INTRODUCTION

A complementary cosmological observations confirm that our universe is currently undergoing a phase of accelerated expansion [1]. Within the framework of general relativity (GR), the acceleration can be explained by assuming a bizarre type of energy with a negative pressure called “dark energy” (DE). A plenty of models have been proposed to explain the acceleration of the universe expansion. One can refer to [2, 3] for a review of DE models. On the other side, many people believe in a modification of gravity, seeking an explanation for the late time acceleration. According to this idea the acceleration will be a part of the universe expansion and does not need to invoke any kind DE component. As examples of this approach one can look at Refs. [4–7].

DE models are mainly based on introducing a new degree of freedom or a new parameter which provide the needed conditions for explanation of the cosmic acceleration (see e.g. [8] and references therein). However, it would be nice to resolve the DE puzzle without presenting any new degree of freedom or any new parameter in the theory. One of the successful and beautiful theories of modern physics is Quantum Chromodynamics (QCD) which describes the strong interaction in nature. However, resolution of one of its mysteries, the U(1) problem, has remained somewhat unsatisfying. Veneziano ghost field explained the U(1) problem in QCD [9]. Vacuum energy of the ghost field can be used to explain the time varying cosmological constant in a spacetime with nontrivial topology, since the ghost filed has no contribution to the vacuum energy in the Minkowskian spacetime [10]. The ghost vacuum energy density is proportional to $\Lambda^3_{QCD}H$, where Λ_{QCD} is QCD mass scale and H is the Hubble parameter [11]. It is well-known that the cosmological constant model of DE suffers the coincidence and the fine tuning problems. However, with correct choice of Λ_{QCD}, the ghost dark energy (GDE) model does not encounter the fine tuning problem anymore [10, 11]. Phenomenological implications of GDE model were discussed in [12]. In [13] the GDE in a non-flat universe in the presence of interaction between DE and dark matter was explored. The instability of GDE model against perturbations was studied in [14]. It was argued that the perfect fluid for GDE is classically unstable against perturbations. Other features of GDE model have been investigated in Refs. [15–17].

In all above references ([10–21]) the GDE was assumed to have the energy density of the form $\rho_D = \alpha H$, while, in general, the vacuum energy of the Veneziano ghost field in QCD is of the form $H + O(H^2)$ [22]. This indicates that in the previous works on the GDE model, only the leading term H has been considered. Motivated by the argument given in [23], one may expect that the subleading term H^2 in the GDE model might play a crucial role in the early evolution of the universe, acting as the early DE. It was shown [24] that taking the second term into account can give better agreement with observational data compared to the usual GDE. Hereafter we call this model as the generalized ghost dark energy (GGDE) and our main task in this paper is to investigate the main properties of this model. In this model the energy density is written in the form $\rho_D = \alpha H + \beta H^2$, where β is a constant.

Beside the DE component in the universe there exist another unknown component called "dark matter" (DM). Since the nature of these two dark components are still mystery and they seem to have different gravitational behavior, people usually consider them separately and take their evolution independent of each other. However, there exist

---

* eebrahimi@uk.ac.ir
† sheykhi@uk.ac.ir

several observational evidences which detect signatures of interaction between the two dark components. For example, observational evidences provided by the galaxy cluster Abell A586 support the interaction between DE and DM [25]. Also, it has been shown that interaction can be compatible with SNIa and CMB data [26]. Considering the interaction can also be understood from theoretical point of view. When we assume that the DM and the DE components evolve separately it means that both of them are conserved independently. Any conservation results from a symmetry in the Lagrangian level and in the absence of such a symmetry it seems acceptable to take an interaction between DE and DM.

Based on the cosmological principle the universe has three distinct geometry, namely open, flat and closed geometry corresponding to $k = -1, 0, +1$, respectively. For a long time it was a general belief that universe has a flat ($k = 0$) geometry which mainly based on the inflation theory [27]. With the development of the observational technics people found deviations from the flat geometry [28]. For example, CMB experiments [29] leave the chance for a spatially closed universe. In addition to CMB, recently the spatial geometry of the universe was probed by supernova measurements of the cubic correction to the luminosity distance [30], where a closed universe is also marginally favored. Also the information resulted from WMAP data provides further confidence, showing that a closed universe with positively curved space is marginally preferred [31].

All above reasons, indicate that although people believe in a flat geometry for the universe, but astronomical observations leave enough room for considering a nonflat geometry. Also about the interaction between DM and DE there are several signals from nature which guides us to let the models explain such a behavior. Based on these motivations we would like here to extend the studies on GGDE, to a nonflat FRW spacetime in the presence of an interaction term.

This paper is organized as follows. In the next section, we review the GGDE model in a flat universe. Section III is devoted to the interacting GGDE in a flat geometry. In section IV, we generalize the study to the universe with spacial curvature in the presence of interaction between DM and DE. We summarize our results in section V.

II. GGDE MODEL IN FLAT UNIVERSE

We start with studying the non interacting GGDE model in a flat universe. Although this model has been studied in [24], but we would like to consider the GGDE in a different manner by direct look at the important parameters such as the EoS parameter, w_D, and the deceleration parameter, q.

For flat Friedmann-Robertson-Walker (FRW) universe, the first Friedmann equation takes the form

$$H^2 = \frac{8\pi G}{3}\left(\rho_m + \rho_D\right), \tag{1}$$

where ρ_m and ρ_D are, respectively, the energy densities of pressureless DM and DE. The generalized ghost energy density is defined as [24]

$$\rho_D = \alpha H + \beta H^2, \tag{2}$$

where α is a constant of order $\Lambda^3_{\rm QCD}$ and $\Lambda_{\rm QCD}$ is QCD mass scale, and β is also a constant. In the original GDE ($\beta = 0$) with $\Lambda_{\rm QCD} \sim 100MeV$ and $H \sim 10^{-33}eV$, $\Lambda^3_{\rm QCD}H$ gives the right order of magnitude $\sim (3\times 10^{-3}{\rm eV})^4$ for the observed DE density [11]. In the GGDE, β is a free parameter and can be adjusted for better agreement with observations.

We define the dimensionless energy density parameters as

$$\Omega_m = \frac{\rho_m}{\rho_{cr}} = \frac{8\pi G\rho_m}{3H^2}, \quad \Omega_D = \frac{\rho_D}{\rho_{cr}} = \frac{8\pi G(\alpha+\beta H)}{3H}, \tag{3}$$

where $\rho_{cr} = 3H^2/(8\pi G)$. According to these definitions, the first Friedmann equation can be rewritten in terms of the energy density parameters as

$$\Omega_m + \Omega_D = 1. \tag{4}$$

Through this section we consider GGDE in the absence of interaction term, thus the conservation equations read

$$\dot\rho_m + 3H\rho_m = 0, \tag{5}$$

$$\dot\rho_D + 3H\rho_D(1+w_D) = 0, \tag{6}$$

which imply that DM and DE are separately conserved.



FIG. 1: These figures show the evolutions of w_D and q against Ω_D in a flat GGDE and GDE models. Solid lines correspond to GGDE when $\xi = 0.1$ and the dashed lines belong to GDE model.

Taking time derivative of relation (2), we obtain

$$\dot{\rho}_D = \dot{H}(\alpha + 2\beta H). \tag{7}$$

Also it can be easily seen that differentiating Eq. (1) with respect to time leads to

$$\dot{H} = -4\pi G\rho_D(1 + u + w_D), \tag{8}$$

where $u = \rho_m/\rho_D$ is the energy densities ratio. Replacing this relation in Eq. (7) and also using the continuity equation (6), we get

$$(1 + w_D)[3H - 4\pi G(\alpha + 2\beta H)] = 4\pi G(\alpha + 2\beta H). \tag{9}$$

Solving the above equation for w_D and noticing that $u = \Omega_m/\Omega_D$ and

$$\frac{4\pi G}{3H}(\alpha + 2\beta H) = \frac{\Omega_D}{2} + \frac{4\pi G\beta}{3}, \tag{10}$$

we obtain

$$w_D = \frac{\xi - \Omega_D}{\Omega_D(2 - \Omega_D - \xi)}, \tag{11}$$

where $\xi = \frac{8\pi G\beta}{3}$. It is clear that this relation reduces to its respective one in the GDE when $\xi = 0$ [13].

In Fig. 1a we plotted the evolution of w_D versus Ω_D. It is easy to see that at the late time where $\Omega_D \to 1$, we have $w_D \to -1$, which implies that the GGDE model mimics a cosmological constant behavior. One should notice that this behavior is the same as for the original GDE model. This is expected since the subleading term H^2 in the late time can be ignored due to the smallness of H and the difference between these two models appears only at the early epoches of the universe. From figure (1a) we see that w_D of the GGDE model cannot cross the phantom divide and the universe has a de Sitter phase at the late time. It is important to note that the universe is filled with two dark components namely, DM and GGDE. Thus to discuss the acceleration of the universe we should define the effective EoS parameter, $w_{\rm eff}$, as

$$w_{\rm eff} = \frac{p_t}{\rho_t} = \frac{p_D}{\rho_D + \rho_m}, \tag{12}$$

where ρ_t and p_t are, respectively, the total energy density and the total pressure of the universe. As usual, we have assumed the DM is in the form of pressureless fluid (p_m=0). Using relation (4) for the spatially flat universe, one can find

$$w_{\rm eff} = \Omega_D w_D = \frac{\xi - \Omega_D}{2 - \Omega_D - \xi}. \tag{13}$$

Let us now turn to the deceleration parameter which is defined as

$$q = -\frac{a\ddot{a}}{\dot{a}^2} = -1 - \frac{\dot{H}}{H^2}, \tag{14}$$

where a is the scale factor. Using Eq. (8) and definition Ω_D in (3) we obtain

$$\frac{\dot{H}}{H^2} = -\frac{3}{2}\Omega_D\left(1 + u + w_D\right). \tag{15}$$

Replacing this relation into (14), and using (11) we find

$$q = \frac{1}{2} - \frac{3}{2}\frac{\xi - \Omega_D}{(\xi + \Omega_D - 2)}. \tag{16}$$

One can easily check that the deceleration parameter in GDE is retrieved for $\xi = 0$ [13]. We can also take a look at the early and the late time behavior for the deceleration parameter of this model. At the early time where $\Omega_D \to 0$ the deceleration parameter becomes

$$q = \frac{1}{2} - \frac{3}{2}\frac{\xi}{\xi - 2}. \tag{17}$$

which indicates that for $\xi < 2$ the universe is at the deceleration phase at early times while for $\xi > 2$, the universe could experience an acceleration phase, the former is consistent with the definition $\xi = \frac{8\pi G\beta}{3}$. On the other side, we find that at the late time where the DE dominates ($\Omega_D \to 1$), independent of the value of the ξ, we have $q = -1$. We have plotted the behavior of q in Fig. 1b. Besides, taking $\Omega_{D0} = 0.72$ and adjusting $\xi = 0.01$ we obtain $q_0 \approx -0.34$ for the present value of the deceleration parameter which is in agreement with recent observational data [32]. Choosing the same set of parameters leads to $w_{D0} \approx -0.78$ and $w_{\mathrm{eff}0} \approx -0.56$. One should note that as we already mentioned about w_D, the square term in the GGDE density has a negative contribution in the role of the DE in the universe. We mean by negative contribution that arises by taking the square term into account, the evolution of the universe will be slowed. For example, the universe will enter the acceleration phase later than the original GDE. This behavior is clearly seen in both parts of Fig. 1.

At the end of this section we present the evolution equation of the DE density parameter Ω_D. To this goal we take time derivative of Eq. (3), after using relation $\dot{\Omega}_D = H\frac{d\Omega_D}{d\ln a}$ as well as Eq. (14) we reach

$$\frac{d\Omega_D}{d\ln a} = -3\Omega_D\left(1 - \Omega_D\right)w_D. \tag{18}$$

Using Eq. (11) we get

$$\frac{d\Omega_D}{d\ln a} = -3\frac{(1 - \Omega_D)(\xi - \Omega_D)}{2 - \Omega_D - \xi}. \tag{19}$$

Once again for the limiting case $\xi = 0$, the above relation reduces to its respective evolution equation for the original GDE presented in [13].

III. INTERACTING GGDE IN FLAT UNIVERSE

In the previous section, the evolution of DE and DM components were discussed separately. Here we would like to extend the study to the interacting case, seeking new features of GGDE. In the first look investigating interacting models of DE are valuable from two perspective. The first is the theoretical one, which states that we have no reason against interaction between DE and DM components. For example, in the unified models of field theory DM and DE can be explained by a single scalar field, thus they will be allowed to interact minimally. Besides, one can get rid of the coincidence problem by taking into account the interaction term between DM and DE. One can refer to[33–37] for detailed discussion. The other feature which motivates us to consider interacting models of DE and DM comes from observations. For instance, observational evidences provide by the galaxy cluster Abell A586 support the interaction between DE and DM [25]. Thus, there exist enough motivations to consider the GGDE in the presence of an interaction term. To this end, we start with the energy balance equations for DE and DM, namely

$$\begin{aligned} \dot{\rho}_m + 3H\rho_m &= Q, \qquad (20)\\ \dot{\rho}_D + 3H\rho_D(1 + w_D) &= -Q, \qquad (21) \end{aligned}$$

where $Q > 0$ represents the interaction term which allows the transition of energy from DE to DM. The form of Q is a matter of choice and can be taken as [13]

$$Q = 3b^2H(\rho_m + \rho_D) = 3b^2H\rho_D(1 + u), \tag{22}$$



FIG. 2: These figures show the evolutions of w_D and q against Ω_D for a flat interacting GGDE and GDE model. Solid lines correspond to the GGDE when $\xi = 0.1$ and the dashed lines belong to the GDE model. For both cases $b = 0.15$.

with b^2 being a coupling constant. Inserting Eqs. (7) and (22) in Eq. (21) and taking into account $u = \frac{\Omega_m}{\Omega_D}$, we find

$$w_D = -\frac{1}{2-\Omega_D-\xi}\left(1+\frac{2b^2}{\Omega_D}-\frac{\xi}{\Omega_D}\right). \tag{23}$$

At first look one can find that setting $b = 0$, w_D reduces to the respective relation in the absence of interaction obtained in Eq. (11). When $\xi = 0$ the result recovers those in [13] for original GDE. The first interesting point about the EoS parameter of the GGDE is that in the interacting case independent of the interaction parameter, b, for $0 < \xi < 1$, w_D can cross the phantom line in the future where $\Omega_D \to 1$. At the present time, by choosing $\xi = 0.03$, $b = 0.15$ and $\Omega_{D0} = 0.72$, we find that $w_{D0} = -0.82$ and $w_{\rm eff0} = -0.59$ which the latter favored by observations. One can easily check that for a same coupling constant these values for the original GDE are $w_{D0} = -0.83$ and $w_{\rm eff0} = -0.60$ which clearly show that the square term in the energy density of the GGDE slow down the evolution of the universe compared to the original GDE model. For a better insight we have plotted w_D against Ω_D in Fig. 2a. This value for coupling constant, b, in the figure is consistent with recent observations [38]. It is worth mentioning that at the late time where $\Omega_D \to 1$ the effective EoS parameter approaches less than -1, i.e. $w_{\rm eff} < -1$, which reminds a super acceleration for the universe in the future. Next we take a look at the deceleration parameter in the presence of an interaction term. Substituting (15) in (14) and using (23) yields

$$q = \frac{1}{2} - \frac{3}{2}\frac{\Omega_D}{(2-\Omega_D-\xi)}\left(1+\frac{2b^2}{\Omega_D}-\frac{\xi}{\Omega_D}\right). \tag{24}$$

Once again it is clear that setting $b = 0$, the respective relation in the previous section is retrieved. When $\xi = 0$ the result of [13] is recovered. For the set of parameters $(\xi = 0.03, b = 0.15, \Omega_{D0} = 0.72)$, we find that according to the GGDE the universe enters the acceleration phase at $\Omega = 0.48$ while this transition happens earlier for the GDE model. This point is clear from Fig.2b. The present value of the deceleration parameter for the interacting GGDE model is $q_0 = -0.38$ which is consistent with observations [32].

Finally, we would like to obtain the evolution equation of DE in the presence of interaction. First we take the time derivative of (3) and obtain

$$\dot{\Omega} = \Omega\left[\frac{\dot{\rho}}{\rho} - 2\frac{\dot{H}}{H}\right]. \tag{25}$$

Using relation (21) as well as (15), it is a matter of calculation to show

$$\frac{d\Omega_D}{d\ln a} = 3\Omega_D\left[\frac{1-\Omega_D}{2-\Omega_D-\xi}\left(1+\frac{2b^2}{\Omega_D}-\frac{\xi}{\Omega_D}\right)-\frac{b^2}{\Omega_D}\right]. \tag{26}$$

In the limiting case $\xi = 0$ the equation of motion of interacting GDE is recovered [13].

IV. INTERACTING GGDE IN NON-FLAT UNIVERSE

Flatness problem in standard cosmology was resolved by considering an inflation phase in the evolution history of the universe. Following this theory it became a general belief that our universe is spatially flat. However, later it was



FIG. 3: These figures show the evolutions of w_D and q against Ω_D for a interacting GGDE and GDE models in a non-flat universe. Solid lines correspond to the GGDE when $\xi = 0.1$ and the dashed lines belong to the GDE model. For both cases $b = 0.15$.

shown that exact flatness is not a necessary consequence of inflation if the number of e-foldings is not very large [39]. So it is still possible that there exist a contribution to the Friedmann equation from the spatial curvature, though much smaller than other energy components according to observations. Thus, theoretically the possibility of a curved FRW background is not rejected. In addition, recent observations support the possibility of a non-flat universe and detect a small deviation from $k = 0$ [40]. For example evidences from CMB and supernova measurements of the cubic correction to the luminosity distance favor a positively curved universe [41, 42]. Besides, some exact analysis of the WMAP data support a closed geometry for our universe [43]. Furthermore, the parameter Ω_k represents the contribution to the total energy density from the spatial curvature and it is constrained as $-0.0175 < \Omega_k < 0.0085$ with 95% confidence level by current observations [44]. Our aim in this section is to study the dynamic evolution of the GGDE in a universe with spatial curvature. As we discussed in the introduction a closed universe is marginally favored. We start with the first Friedmann equation in a non-flat universe which is written as

$$H^2 + \frac{k}{a^2} = \frac{1}{3M_p^2}\left(\rho_m + \rho_D\right), \tag{27}$$

where k is the curvature parameter with $k = -1, 0, 1$ corresponding to open, flat, and closed universes, respectively. Taking the energy density parameters (3) into account and defining the energy density parameter for the curvature term as $\Omega_k = k/(a^2H^2)$, the Friedmann equation can be rewritten in the following form

$$1 + \Omega_k = \Omega_m + \Omega_D. \tag{28}$$

Using the above equation the energy density ratio becomes

$$u = \frac{\rho_m}{\rho_D} = \frac{\Omega_m}{\Omega_D} = \frac{1 + \Omega_k - \Omega_D}{\Omega_D}. \tag{29}$$

The second Friedmann equation reads

$$\dot{H} = -4\pi G(p + \rho) + \frac{k}{a^2}, \tag{30}$$

while the time derivative of GGDE density is

$$\dot{\rho}_D = \dot{H}(\alpha + 2\beta H). \tag{31}$$

Inserting Eq. (30) into (31) and combining resulting relation with conservation equation for DE component (21), after using (22) and (29), we find the EoS parameter of interacting GGDE in non-flat universe

$$w_D = -\frac{1}{2 - \Omega_D - \xi}\left(2 - \left(1 + \frac{\xi}{\Omega_D}\right)\left(1 + \frac{\Omega_k}{3}\right) + \frac{2b^2}{\Omega_D}(1 + \Omega_k)\right). \tag{32}$$

From the second Friedmann equation, (30), one can easily obtain

$$\frac{\dot{H}}{H^2} = -\Omega_k + \frac{3}{2}\Omega_D[1 + u + w_D], \tag{33}$$

and therefore the deceleration parameter in a non-flat background is obtained as

$$q = -1 - \frac{\dot{H}}{H^2} = -1 - \Omega_k + \frac{3}{2}\Omega_D[1 + u + w_D]. \tag{34}$$

Substituting Eqs. (29) and (32) in (34) we obtain

$$q = \frac{1}{2}\left(1 + \Omega_k\right) - \frac{3\Omega_D}{2(2 - \Omega_D - \xi)}\left[2 - \left(1 + \frac{\xi}{\Omega_D}\right)\left(1 + \frac{\Omega_k}{3}\right) + \frac{2b^2}{\Omega_D}(1 + \Omega_k)\right]. \tag{35}$$

In a non-flat FRW universe, the equation of motion of interacting GGDE is obtained following the method of the previous section. The result is

$$\frac{d\Omega_D}{d\ln a} = 3\Omega_D\left[\frac{\Omega_k}{3} + \frac{1 - \Omega_D}{2 - \Omega_D - \xi}\left(2 - \left(1 + \frac{\xi}{\Omega_D}\right)\left(1 + \frac{\Omega_k}{3}\right) + \frac{2b^2}{\Omega_D}(1 + \Omega_k)\right) - \frac{b^2}{\Omega_D}(1 + \Omega_k)\right]. \tag{36}$$

In the limiting case $\Omega_k = 0$, the results of this section restore their respective equations in flat FRW universe derived in the previous sections, while for $\xi = 0$ the respective relations in [13] are retrieved. The evolutions of w_D and q against Ω_D for a non-flat interacting GGDE and GDE models are plotted in Fig.3. Let us explore different features of GGDE in non-flat universe by a numerical study. First of all we study the EoS parameter of the GGDE in the future where $\Omega_D \to 1$. In this case, taking $\xi = 0.1$, $b = 0.15$ and $\Omega_k = 0.01$ leads to $w_D = -1.05$ which indicates that the GGDE is capable to cross the phantom line in the future. The present stage of the universe can be achieved by the same set of parameters but $\Omega_D = 0.72$. In such a case we see that $w_{D0} = -0.78$ while the effective EoS parameter becomes $w_{\mathrm{eff}0} = -0.6$ which is consistent with observations. The deceleration parameter of the model can also be obtained which is in agreement with observational evidences. For example, for the above choice of parameters one finds $q_0 = -0.34$ [32]. Transition from deceleration to the acceleration phase, in the interacting non-flat case, take place at $\Omega_D = 0.52$.

V. SUMMARY AND DISCUSSION

Veneziano ghost model was presented to solve U(1) problem in QCD theory. People usually prefer to handle the DE problem in physics using degree's of freedom already exist. GDE is a prototype of these models which discusses the acceleration of the universe and is based on the mentioned Veneziano ghost field. This model of DE can address the fine tuning problem [13]. An extended version of this model called GGDE was recently proposed by Cai et. al., [24], seeking a better agreement with observations.

In this paper we explored some features of GGDE in both flat and non-flat FRW universe in the presence of an interaction term between the two dark components of the universe. In section II, we discussed the GGDE in a flat FRW background. We found that the EoS parameter approaches -1 which is the same as the cosmological constant. The next section was devoted to the interacting GGDE in flat geometry. An interesting feature which we found was the capability of crossing the phantom line in this case. This behavior is also seen in the last section for interacting GGDE in a universe with spatial curvature.

We also investigated the deceleration parameter in different cases. For flat universe in the presence of interaction between DE and DM we found that with $\Omega_D = 0.72$ for present time and choosing $\xi = 0.01$ and $b = 0.15$, we reach $q = -0.38$ which shows a consistent value in comparison with observations [32]. Also this parameter has a consistent behavior in all cases of interest.

In summary the main result found through this paper is that in the GGDE model, there is a delay in different epoches of the cosmic evolution in comparison with original GDE model. For instance we saw that the deceleration parameter for the present time in the flat noninteracting GDE was -0.34 while for GGDE, the same parameter was obtained -0.33 for $\xi = 0.03$. This point is completely seen from the figures presented in the paper. This result were also pointed in [24] due to the negative contribution of the square term in the energy density of GGDE.

Acknowledgments

This work has been supported financially by Research Institute for Astronomy and Astrophysics of Maragha (RIAAM) under research project number No. 1/2782-62.

[1] A.G. Riess, et al., Astron. J. 116 (1998) 1009;
S. Perlmutter, et al., Astrophys. J. 517 (1999) 565;
S. Perlmutter, et al., Astrophys. J. 598 (2003) 102;
P. de Bernardis, et al., Nature 404 (2000) 955.

[2] E. J. Copeland, M. Sami and S. Tsujikawa, Int. J. Mod. Phys. D 15, 1753 (2006).

[3] M. Li, X. -D. Li, S. Wang, Y. Wang, Commun. Theor. Phys. 56, 525-604 (2011).

[4] G. R. Dvali, G. Gabadadze and M. Porrati, Phys. Lett. B 485 (2000) 208.

[5] C. Deffayet, Phys. Lett. B 502 (2001) 199.

[6] N. Arkani Hamed, H. C. Cheng, M. A. Luty and S. Mukohyama, JHEP 05 (2004) 074.

[7] N. Arkani Hamed, H. C. Cheng, M.A. Luty, S. Mukohyama and T. Wiseman, JHEP 01 (2007) 036.

[8] A. Sheykhi, Phys. Lett. B 680 (2009) 113;
A. Sheykhi, Class. Quantum Gravit. 27 (2010) 025007;
A. Sheykhi, Phys. Lett. B681 (2009) 205;
K. Karami, et. al., Gen. Relativ. Gravit. 43 (2011) 27;
M. Jamil, A. Sheykhi, Int. J. Theor. Phys. 50 (2011) 625;
A. Sheykhi, M. Jamil, Phys. Lett. B 694 (2011) 284.

[9] E. Witten, Nucl. Phys. B 156 (1979) 269;
G. Veneziano, Nucl. Phys. B 159 (1979) 213;
C. Rosenzweig, J. Schechter and C. G. Trahern, Phys. Rev. D 21 (1980) 3388.

[10] F. R. Urban and A. R. Zhitnitsky, Phys. Lett. B 688 (2010) 9;
K. Kawarabayashi and N. Ohta, Nucl. Phys. B 175 (1980) 477.

[11] N. Ohta, Phys. Lett. B 695 (2011) 41, arXiv:1010.1339.

[12] R.G. Cai, Z.L. Tuo, H.B. Zhang, arXiv:1011.3212.

[13] A. Sheykhi, M.Sadegh Movahed, Gen Relativ Gravit 44 (2012) 449.

[14] E. Ebrahimi and A. Sheykhi, Int. J. Mod. Phys. D20 (2011) 2369.

[15] A. Sheykhi, M. Sadegh Movahed, E. Ebrahimi, Astrophys Space Sci 339 (2012)93.

[16] A. Sheykhi, A. Bagheri, Euro. Phys. Lett., 95 (2011) 39001.

[17] E. Ebrahimi and A. Sheykhi, Phys. Lett. B 706 (2011) 19.

[18] A. Rozas-Fernandez, Phys.Lett. B709 (2012) 313-321.

[19] K. Karami, M. Mousivand, arXiv:1209.2044.

[20] A. Khodam-Mohammadi, M. Malekjani, M. Monshizadeh, Mod. Phys. Lett. A27, 18 (2012) 1250100.

[21] M. Malekjani, A. Khodam-Mohammadi, arXiv:1202.4154.

[22] A. R. Zhitnitsky, arXiv:1112.3365.

[23] M. Maggiore, L. Hollenstein, M. Jaccard and E. Mitsou, Phys. Lett. B 704, 102 (2011).

[24] R. G. Cai, Z. L. Tuo, Y. B. Wu, Y. Y. Zhao, arXiv:1201.2494

[25] O. Bertolami , F. Gil Pedro and M. Le Delliou, Phys. Lett. B 654 (2007) 165.

[26] G. Olivares, F. Atrio, D. Pavon, Phys. Rev. D 71 (2005) 063523.

[27] A. H. Guth, Phys. Rev. D 23,347 (1981).

[28] C. L. Bennett, et al., Astrophys. J. Suppl. 148 (2003) 1;
D. N. Spergel, Astrophys. J. Suppl. 148 (2003) 175;
M. Tegmark, et al., Phys. Rev. D 69 (2004) 103501;
U. Seljak, A. Slosar, P. McDonald, JCAP 0610 (2006) 014;
D. N. Spergel, et al., Astrophys. J. Suppl. 170 (2007) 377.

[29] J. L. Sievers, et al., Astrophys. J. 591 (2003) 599;
C.B. Netterfield, et al., Astrophys. J. 571 (2002) 604;
A. Benoit, et al., Astron. Astrophys. 399 (2003) L25;
A. Benoit, et al., Astron. Astrophys. 399 (2003) L19.

[30] R. R. Caldwell, M. Kamionkowski, astro-ph/0403003;
B. Wang, Y. G. Gong, R. K. Su, Phys. Lett. B 605 (2005) 9.

[31] J. P. Uzan, U. Kirchner, G.F.R. Ellis, Mon. Not. R. Astron. Soc. 344 (2003) L65;
A. Linde, JCAP 0305 (2003) 002;
M. Tegmark, A. de Oliveira-Costa, A. Hamilton, Phys. Rev. D 68 (2003) 123523;
G. Efstathiou, Mon. Not. R. Astron. Soc. 343 (2003) L95;
J. P. Luminet, J. Weeks, A. Riazuelo, R. Lehou, J. Uzan, Nature 425 (2003) 593;
G. F. R. Ellis, R. Maartens, Class. Quantum Grav. 21 (2004) 223.

[32] R.A. Daly, et al., J. Astrophys. 677 (2008) 1.
[33] L. Amendola, Phys. Rev. D 62, 043511 (2000);
L. Amen- dola and C. Quercellini, Phys. Rev. D 68 (2003) 023514 ;
L. Amendola, S. Tsujikawa and M. Sami, Phys. Lett. B 632 (2006) 155.
[34] D. Pavon,W. Zimdahl, Phys. Lett. B 628 (2005) 206;
S. Campo, R. Herrera, D. Pavon, Phys. Rev. D 78 (2008) 021302(R).
[35] C. G. Boehmer, G. Caldera-Cabral, R. Lazkoz, R. Maartens, Phys. Rev. D 78 (2008) 023505.
[36] G. Olivares, F. Atrio-Barandela and D. Pavon, Phys. Rev. D 74 (2006) 043521.
[37] S. B. Chen, B. Wang, J. L. Jing, Phys.Rev. D 78 (2008) 123503.
[38] B. Wang, Y. Gong and E. Abdalla, Phys. Lett. B 624 (2005) 141;
B. Wang, C. Y. Lin and E. Abdalla, Phys. Lett. B 637 (2005) 357.
[39] Q. G. Huang and M. Li, J. Cosmol. Astropart. Phys. JCAP08 (2004)013.
[40] C. L. Bennett, et al., Astrophys. J. Suppl. 148 (2003) 1;
D. N. Spergel, Astrophys. J. Suppl. 148 (2003) 175;
M. Tegmark, et al., Phys. Rev. D 69 (2004) 103501;
U. Seljak, A. Slosar, P. McDonald, JCAP 0610 (2006) 014;
D. N. Spergel, et al., Astrophys. J. Suppl. 170 (2007) 377.
[41] J. L. Sievers, et al., Astrophys. J. 591 (2003) 599;
C.B. Netterfield, et al., Astrophys. J. 571 (2002) 604;
A. Benoit, et al., Astron. Astrophys. 399 (2003) L25;
A. Benoit, et al., Astron. Astrophys. 399 (2003) L19.
[42] R. R. Caldwell, M. Kamionkowski, JCAP 0409 (2004) 009;
B. Wang, Y. G. Gong, R. K. Su, Phys. Lett. B 605 (2005) 9.
[43] J. P. Uzan, U. Kirchner, G.F.R. Ellis, Mon. Not. R. Astron. Soc. 344 (2003) L65;
A. Linde, JCAP 0305 (2003) 002;
M. Tegmark, A. de Oliveira-Costa, A. Hamilton, Phys. Rev. D 68 (2003) 123523;
G. Efstathiou, Mon. Not. R. Astron. Soc. 343 (2003) L95;
J. P. Luminet, J. Weeks, A. Riazuelo, R. Lehou, J. Uzan, Nature 425 (2003) 593;
G. F. R. Ellis, R. Maartens, Class. Quantum Grav. 21 (2004) 223.
[44] T. P. Waterhouse and J. P. Zipin, arXiv:0804.1771.